



SHINGLES (HERPES ZOSTER)

Questions and answers

► What is shingles?

Shingles, also called herpes zoster or zoster, is a painful skin rash caused by the varicella zoster virus (VZV). VZV is the same virus that causes chickenpox. After a person recovers from chickenpox, the virus stays in the body. Usually the virus does not cause any problems; however, the virus can reappear years later, causing shingles. Herpes zoster is not caused by the same virus that causes genital herpes, a sexually transmitted disease.

► What does shingles look like?

Shingles usually starts as a rash on one side of the face or body. The rash starts as blisters that scab after three to five days. The rash usually clears within two to four weeks. Before the rash develops, there is often pain, itching or tingling in the area where the rash will develop. Other symptoms of shingles can include fever, headache, chills and upset stomach.

► Are there any long-term effects from shingles?

Very rarely, shingles can lead to pneumonia, hearing problems, blindness, brain inflammation (encephalitis) or death. For about one person in five, severe pain can continue even after the rash clears up. This pain is called post-herpetic neuralgia. As people get older, they are more likely to develop post-herpetic neuralgia and it is more likely to be severe.

► How common is shingles in the United States?

In the United States, there are an estimated one million cases of shingles each year. About one of every three people in the United States will get shingles during their lifetime. Approximately half of those who live to be 85 years old will have one or more encounters with shingles.

► Who gets shingles?

Anyone who has recovered from chickenpox may develop shingles, including children. However, shingles most commonly occurs in people 50 years old and older. The risk of getting shingles increases as a person gets older:

People who have medical conditions that keep the immune system from working properly, like cancer; leukemia, lymphoma and human immunodeficiency virus (HIV), or people who receive immunosuppressive drugs, such as steroids and drugs given after organ transplantation, are also at greater risk to get shingles.

► How often can a person get shingles?

Most commonly, a person has only one episode of shingles in their lifetime. Although rare, a second or even third case of shingles can occur.

► Can shingles be spread to others?

Shingles cannot be passed from one person to another. However, the virus that causes shingles, VZV, can be spread from a person with active shingles to a person who has never had chickenpox through direct contact with the rash. The person exposed would develop chickenpox, not shingles.

The virus is not spread through sneezing, coughing or casual contact. A person with shingles can spread the disease when the rash is in the blister phase. Once the rash has developed crusts, the person is no longer contagious.

A person is not infectious before blisters appear or with post-herpetic neuralgia (pain after the rash is gone).

Continued on back

SHINGLES (HERPES ZOSTER): *Questions and answers*

► What can be done to prevent the spread of shingles?

The risk of spreading shingles is low if the rash is covered. People with shingles should keep the rash covered, not touch or scratch the rash, and wash their hands often to prevent the spread of VZV. Once the rash has developed crusts, the person is no longer contagious.

► Is there a treatment for shingles?

Several medicines, acyclovir (Zovirax), valacyclovir (Valtrex) and famciclovir (Famvir) are available to treat shingles. These medications should be started as soon as possible after the rash appears and will help shorten how long the illness lasts and how severe the illness is. Pain medicine may also help with pain caused by shingles. Call your doctor as soon as possible to discuss treatment options.

► Is there a vaccine to prevent shingles?

Yes. There are two vaccines that prevent shingles. One for use in people ages 60 and older which is a single dose and another for ages 50 and older that requires two doses to prevent shingles. Neither vaccine is used to treat shingles or post-herpetic neuralgia (pain after the rash is gone) once it develops.

► Does the shingles vaccine work?

Yes. Both vaccines work. Shingrix protected almost 100% of people from getting the shingles rash and about 90% from the long-lasting pain associated with shingles. Zostavax protected more than half of the recipients from getting the shingles rash and about two-thirds from getting long-lasting shingles pain. Because of the dramatic difference between these two vaccines, the committee of experts that advises the Centers for Disease Control and Prevention (CDC) recommended a preference for Shingrix vaccine.

► Can the shingles vaccine be given to people who have already had shingles?

Yes. People who have had shingles can receive the shingles vaccine to help prevent future occurrences of the disease.

► Why is the shingles vaccine only recommended for people 50 years and older?

A person's risk for getting shingles begins to rise around age 50.

► How are vaccine recommendations made?

Once a vaccine is licensed by the FDA, the federal Advisory Committee on Immunization Practices (ACIP) votes on whether to recommend this vaccine, and if so, who should get it and at what ages. Neither ACIP nor the federal government

makes mandates or laws requiring immunization for adults. Recommendations made by ACIP will be reviewed by the CDC director and the Department of Health and Human Services. Recommendations become official when published in the CDC's Morbidity and Mortality Weekly Report (MMWR).

► Has ACIP recommended the FDA-approved vaccine?

Yes. The shingles vaccine was recently recommended by ACIP to reduce the risk of shingles and its associated pain in people 60 years old or older. See MMWR: Prevention of Shingles (ACIP Recommendations).

► Is the FDA-approved vaccine safe?

The FDA has licensed the vaccine as safe. The vaccine has been tested in about 20,000 people ages 60 and older. The most common side effects were redness, soreness, swelling or itching at the shot site, and headache. The CDC, working with the FDA, will continue to monitor the safety of the vaccine after it is in general use.

► Will zoster vaccine be covered by Medicare for Medicare beneficiaries?

While details are evolving, it is anticipated that zoster vaccine will not be covered under Medicare Part B (which covers influenza and pneumococcal polysaccharide vaccine as well as hepatitis B for moderate and high risk people). The vaccine will instead be reimbursed through the Medicare Part D program. Beneficiaries should contact their Part D plan for more information.

For More Information:

■ Louisiana Department of Health Immunization Program
1450 Poydras Street, Suite 1938
New Orleans, LA 70112
Phone: (504) 568-2600
Fax: (504) 568-2659

► Where can I get more information?

- Your doctor, nurse or clinic, a community pharmacist or your local public health unit
- Centers for Disease Control and Prevention (CDC): cdc.gov/shingles/about/index.html

This fact sheet is for information only and is not meant to be used for self-diagnosis or as a substitute for consultation with a health care provider. If you have any questions about the disease described above or think that you may have this infection, consult a health care provider.

